HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use SUTAB $^{\circ}$ safely and effectively. See full prescribing information for SUTAB.

SUTAB (sodium sulfate, magnesium sulfate, and potassium chloride) tablets, for oral use Initial U.S. Approval: 2020

RECENT MAJOR CHANGES			
Contraindications (4, 6.2)			
Warnings and Precautions (5.5)			
Postmarketing Experience (6.2)			

------**DOSAGE AND ADMINISTRATION**-----For complete information on preparation before colonoscopy and administration of the dosage regimen, see full prescribing information. (2.1, 2.2)

- Preparation and Administration (2.1)
- Administration of two doses of SUTAB (24 tablets) are required for a complete preparation for colonoscopy. Twelve (12) tablets are equivalent to one dose.
- Water must be consumed with each dose of SUTAB and additional water must be consumed after each dose.
- Do not take other laxatives while taking SUTAB.
- Do not take oral medications within 1 hour of starting each dose of SUTAB.
- If taking tetracycline or fluoroquinolone antibiotics, iron, digoxin, chlorpromazine, or penicillamine, take these medications at least 2 hours before and not less than 6 hours after administration of each dose of SUTAB.

Split Dose (2-Day) Regimen (2.2)

- Evening before colonoscopy:
 - Open 1 bottle of 12 tablets.
 - Fill the provided container with 16 ounces of water (up to the fill line). Swallow each tablet with a sip of water and drink the entire amount over 15 to 20 minutes.
 - Approximately one hour after the last tablet is ingested, fill the provided container a second time with 16 ounces of water (up to the fill line) and drink the entire amount over 30 minutes.
 - Approximately 30 minutes after finishing the second container of water, fill the provided container with 16 ounces of water (up to the fill line) and drink the entire amount over 30 minutes.
- Morning of the colonoscopy:
 - Open 1 bottle of 12 tablets.
 - Fill the provided container with 16 ounces of water (up to the fill line). Swallow each tablet with a sip of water and drink the entire amount over 15 to 20 minutes.
 - Approximately one hour after the last tablet is ingested, fill the provided container a second time with 16 ounces of water (up to the fill line) and drink the entire amount over 30 minutes.

- Approximately 30 minutes after finishing the second container of water, fill the provided container with 16 ounces of water (up to the fill line) and drink the entire amount over 30 minutes.
- Complete all SUTAB tablets and required water at least 2 hours before colonoscopy.

-----DOSAGE FORMS AND STRENGTHS------DOSAGE FORMS AND STRENGTHS------

Tablets: 1.479 g sodium sulfate, 0.225 g magnesium sulfate, and 0.188 g potassium chloride. (3)

-----CONTRAINDICATIONS------

- Gastrointestinal obstruction or ileus (4, 5.6)
- Bowel perforation (4, 5.6)
- Toxic colitis or toxic megacolon (4)
- Gastric retention (4)
- Hypersensitivity to any ingredient in SUTAB (4, 6.2)

------WARNINGS AND PRECAUTIONS------

- <u>Risk of fluid and electrolyte abnormalities</u>: Encourage adequate hydration, assess concurrent medications and consider laboratory assessments prior to and after each use. (5.1, 7.1)
- <u>Cardiac arrhythmias</u>: Consider pre-dose and post-colonoscopy ECGs in patients at increased risk. (5.2)
- <u>Seizures</u>: Use caution in patients with a history of seizures and patients at increased risk of seizures, including medications that lower the seizure threshold. (5.3, 7.1)
- <u>Patients with renal impairment or taking concomitant medications that affect renal function</u>: Use caution, ensure adequate hydration and consider laboratory testing. (5.4, 7.1)
- <u>Mucosal ulcerations</u>: Consider potential for mucosal ulcerations when interpreting colonoscopy findings in patients with known or suspected inflammatory bowel disease. Consider potential for mucosal ulcerations in patients undergoing esophageal gastroduodenoscopy. (5.5, 6.2)
- <u>Suspected GI obstruction or perforation</u>: Rule out the diagnosis before administration. (4, 5.6)

-----ADVERSE REACTIONS------

Most common gastrointestinal adverse reactions are nausea, abdominal distension, vomiting and upper abdominal pain. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Braintree Laboratories, Inc. at 1-800-874-6756 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

------DRUG INTERACTIONS-------Drugs that increase risk of fluid and electrolyte imbalance. (7.1)

See 17 for PATIENT COUNSELING INFORMATION and Medication Guide.

Revised: 08/2021

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

- 2.1 Preparation and Administration Instructions
- 2.2 Split-Dose (2-Day) Regimen

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

- 5 WARNINGS AND PRECAUTIONS
 - 5.1 Serious Fluid and Electrolyte Abnormalities
 - 5.2 Cardiac Arrhythmias
 - 5.3 Seizures
 - 5.4 Use in Patients with Risk of Renal Injury
 - 5.5 Mucosal Ulcerations and Ischemic Colitis
 - 5.6 Use in Patients with Significant Gastrointestinal Disease

6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing Experience
- 7 DRUG INTERACTIONS

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Lactation
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Renal Impairment
- 10 OVERDOSAGE
- 11 DESCRIPTION
- 12 CLINICAL PHARMACOLOGY
 - 12.1 Mechanism of Action
 - 12.3 Pharmacokinetics
- 13 NONCLINICAL TOXICOLOGY
 - 13.2 Animal Toxicology and/or Pharmacology
- 14 CLINICAL STUDIES
- 16 HOW SUPPLIED/STORAGE AND HANDLING
- 17 PATIENT COUNSELING INFORMATION

- 7.1 Drugs That May Increase Risks of Fluid and Electrolyte Abnormalities
- 7.2 Potential for Reduced Drug Absorption
- 7.3 Stimulant Laxatives

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

SUTAB is indicated for the cleansing of the colon as a preparation for colonoscopy in adults.

2 DOSAGE AND ADMINISTRATION

2.1 Preparation and Administration Instructions

- Correct fluid and electrolyte abnormalities before treatment with SUTAB [see Warnings and Precautions (5.1)]
- Administration of two doses of SUTAB (24 tablets) are required for a complete preparation for colonoscopy. Twelve (12) tablets are equivalent to one dose.
- Water must be consumed with each dose of SUTAB and additional water must be consumed after each dose.
- Consume a low residue breakfast on the day before colonoscopy, followed by clear liquids up to 2 hours prior to colonoscopy.
- Do not drink milk or eat or drink anything colored red or purple.
- Do not drink alcohol.
- Do not take other laxatives while taking SUTAB.
- Do not take oral medications within 1 hour of starting each dose of SUTAB
- If taking tetracycline or fluoroquinolone antibiotics, iron, digoxin, chlorpromazine, or penicillamine, take these
 medications at least 2 hours before and not less than 6 hours after administration of each dose of SUTAB.
- Stop consumption of all fluids at least 2 hours prior to the colonoscopy.

2.2 Split-Dose (2-Day) Regimen

The recommended Split-Dose regimen for adults consists of two doses of SUTAB: the first dose during the evening prior to colonoscopy and the second dose the next day, during the morning of the colonoscopy. Instruct patients:

Dose 1 – On the day prior to colonoscopy:

- A low residue breakfast may be consumed. Examples of low residue foods are eggs, white bread, cottage cheese, yogurt, grits, coffee, tea.
- After breakfast, only clear liquids may be consumed until after the colonoscopy.
- Early in the evening prior to colonoscopy, open one bottle of 12 tablets.
- Fill the provided container with 16 ounces of water (up to the fill line). Swallow each tablet with a sip of water and drink the entire amount over 15 to 20 minutes.
- Approximately one hour after the last tablet is ingested, fill the provided container a second time with 16 ounces
 of water (up to the fill line) and drink the entire amount over 30 minutes.
- Approximately 30 minutes after finishing the second container of water, fill the provided container again with 16 ounces of water (up to the fill line) and drink the entire amount over 30 minutes.
- If patients experience preparation-related symptoms (e.g. nausea, bloating, cramping), pause or slow the rate of drinking the additional water until symptoms diminish.

Dose 2 - Day of colonoscopy:

- Continue to consume only clear liquids until after the colonoscopy.
- The morning of colonoscopy (5 to 8 hours prior to the colonoscopy and no sooner than 4 hours from starting Dose 1), open the second bottle of 12 tablets.
- Fill the provided container with 16 ounces of water (up to the fill line). Swallow each tablet with a sip of water and drink the entire amount over 15 to 20 minutes.
- Approximately one hour after the last tablet is ingested, fill the provided container a second time with 16 ounces
 of water (up to the fill line) and drink the entire amount over 30 minutes.
- Approximately 30 minutes after finishing the second container of water, fill the provided container again with 16 ounces of water (up to the fill line) and drink the entire amount over 30 minutes.
- If patients experience preparation-related symptoms (e.g. nausea, bloating, cramping), pause or slow the rate of drinking the additional water until symptoms diminish.
- Complete all SUTAB tablets and water at least two hours prior to colonoscopy.

3 DOSAGE FORMS AND STRENGTHS

Tablets: 1.479 g sodium sulfate, 0.225 g magnesium sulfate, and 0.188 g potassium chloride. The tablets are white to off-white, film coated, oblong, and biconvex with flat sides, debossed with S24 on one side.

4 CONTRAINDICATIONS

SUTAB is contraindicated in the following conditions:

- Gastrointestinal obstruction or ileus [see Warnings and Precautions (5.6)]
- Bowel perforation [see Warnings and Precautions (5.6)]
- Toxic colitis or toxic megacolon
- Gastric retention
- Hypersensitivity to any ingredient in SUTAB

5 WARNINGS AND PRECAUTIONS

5.1 Serious Fluid and Electrolyte Abnormalities

Advise all patients to hydrate adequately before, during, and after the use of SUTAB. If a patient develops significant vomiting or signs of dehydration after taking SUTAB, consider performing post-colonoscopy lab tests (electrolytes, creatinine, and BUN). Fluid and electrolyte disturbances can lead to serious adverse events including cardiac arrhythmias, seizures and renal impairment. Correct fluid and electrolyte abnormalities before treatment with SUTAB. Use SUTAB with caution in patients with conditions, or who are using medications, that increase the risk of fluid and electrolyte disturbances or may increase the risk of adverse events of seizure, arrhythmias, and renal impairment. *[see Drug Interactions (7.1)]*

- Colonic Mucosal Ulceration and Ischemic Colitis [see Warnings and Precautions (5.5)]
- Patients with Significant Gastrointestinal Disease [see Warnings and Precautions (5.6)]

6.1 Clinical Trials Experience

Because clinical studies are conducted under widely varying conditions, adverse reaction rates observed in the clinical studies of a drug cannot be directly compared to rates in clinical studies of another drug and may not reflect the rates observed in practice.

The safety of SUTAB was evaluated in two randomized, parallel group, multicenter, investigator-blinded clinical trials in 941 adult patients undergoing colonoscopy. The active comparators were polytheylene glycol 3350, sodium sulfate, sodium chloride, potassium chloride, ascorbic acid and sodium ascorbate for oral solution in Study 1 and sodium picosulfate, magnesium oxide, and anhydrous citric acid for oral solution in Study 2 *[see Clinical Studies (14)]*.

Adverse Gastrointestinal Reactions Reported by Symptom Questionnaire

In Studies 1 and 2, patients were queried for selected gastrointestinal adverse reactions of stomach cramping (upper abdominal pain), stomach bloating (abdominal distention), nausea and vomiting using a standard questionnaire following completion of study drug and prior to colonoscopy on the day of colonoscopy. Patients reporting selected gastrointestinal symptom(s) rated the intensity as mild, moderate or severe.

A total of 52% (287/552) of patients in Study 1 and 52% (202/389) in Study 2 reported at least one selected gastrointestinal adverse reaction when queried using the standard questionnaire. Tables 1 and 2 show results for each gastrointestinal adverse reaction reported by patients using the standard questionnaire, including severity.

Table 1: Gastrointestinal Symptoms by Severity^a From Symptom Questionnaire in Adult Patients Following Colon Cleansing and Prior to Colonoscopy – Study 1^b

Symptom	SUTAB	Polyethylene glycol 3350, sodium sulfate, sodium chloride, potassium chloride, ascorbic acid and sodium ascorbate
Total Number of Patients per Treatment Arm (N)	281	271
Patients with at Least One Gastrointestinal Adverse Reaction from Symptom Questionnaire	163	124
% Nausea ^c	48	26
Mild	71	77
Moderate	27	23
Severe	2	0
% Abdominal Distension ^{c,d}	29	22
Mild	68	71
Moderate	30	29
Severe	1	0
% Vomiting ^c	23	5
Mild	48	46
Moderate	52	54
Severe	0	0
% Upper Abdominal Pain ^c	16	18
Mild	65	71
Moderate	35	29
Severe	0	0

^a Mild: barely noticeable, does not influence functioning causing no limitations of usual activities;

Moderate: makes participant uncomfortable, influences functioning causing some limitations of usual activities;

Severe: severe discomfort, treatment needed, severe and undesirable, causing inability to carry out usual activities

^b Study 1 was not designed to support comparative claims for SUTAB for the adverse reactions reported in this table.

^c Percentage represents n/N for patients who experienced each gastrointestinal adverse reaction on the symptom questionnaire based on the total number of patients per treatment arm.

Table 2: Gastrointestinal Symptoms by Severity^a From Symptom Questionnaire in Adult Patients Following Colon Cleansing and Prior to Colonoscopy – Study 2^b

Symptom	SUTAB	Sodium picosulfate, magnesium oxide, and anhydrous citric acid
Total Number of Patients per Treatment Arm (N)	190	199
Patients with at Least One Gastrointestinal Adverse Reaction from Symptom Questionnaire	135	67
% Nausea ^c	52	18
Mild	74	94
Moderate	20	6
Severe	6	0
% Abdominal Distension ^c	34	15
Mild	73	69
Moderate	27	31
Severe	0	0
% Vomiting ^c	16	2
Mild	53	33
Moderate	47	67
Severe	0	0
% Upper Abdominal Pain ^c	23	13
Mild	82	100
Moderate	16	0
Severe	2	0

5.2 Cardiac Arrhythmias

There have been rare reports of serious arrhythmias associated with the use of ionic osmotic laxative products for bowel preparation. Use caution when prescribing SUTAB for patients at increased risk of arrhythmias (e.g., patients with a history of prolonged QT, uncontrolled arrhythmias, recent myocardial infarction, unstable angina, congestive heart failure, or cardiomyopathy). Consider pre-dose and post-colonoscopy ECGs in patients at increased risk of serious cardiac arrhythmias.

5.3 Seizures

There have been reports of generalized tonic-clonic seizures and/or loss of consciousness associated with use of bowel preparation products in patients with no prior history of seizures. The seizure cases were associated with electrolyte abnormalities (e.g., hyponatremia, hypokalemia, hypocalcemia, and hypomagnesemia) and low serum osmolality. The neurologic abnormalities resolved with correction of fluid and electrolyte abnormalities.

Use caution when prescribing SUTAB for patients with a history of seizures and in patients at increased risk of seizure, such as patients taking medications that lower the seizure threshold (e.g., tricyclic antidepressants), patients withdrawing from alcohol or benzodiazepines, or patients with known or suspected hyponatremia [see Drug Interactions (7.1)].

5.4 Use in Patients with Risk of Renal Injury

Use SUTAB with caution in patients with impaired renal function or patients taking concomitant medications that may affect renal function (such as diuretics, angiotensin converting enzyme inhibitors, angiotensin receptor blockers, or non-steroidal anti-inflammatory drugs) *[see Drug Interactions (7.1)]*. These patients may be at risk for renal injury. Advise these patients of the importance of adequate hydration with SUTAB and consider performing baseline and post-colonoscopy laboratory tests (electrolytes, creatinine, and BUN) in these patients *[see Use in Specific Populations (8)]*.

5.5 Mucosal Ulcerations and Ischemic Colitis

Osmotic laxative products may produce colonic mucosal aphthous ulcerations, and there have been reports of more serious cases of ischemic colitis requiring hospitalization. In addition, gastric ulcerations or gastritis may occur. Concurrent use of stimulant laxatives and SUTAB may increase these risks *[see Drug Interactions (7.3)]*. Consider the potential for mucosal ulcerations resulting from the bowel preparation when interpreting colonoscopy findings in patients with known or suspect inflammatory bowel disease (IBD). Consider the potential for mucosal ulcerations resulting from the bowel preparation in patients undergoing esophageal gastroduodenoscopy.

5.6 Use in Patients with Significant Gastrointestinal Disease

If gastrointestinal obstruction or perforation is suspected, perform appropriate diagnostic studies to rule out these conditions before administering SUTAB [see Contraindications (4)].

Use with caution in patients with severe active ulcerative colitis.

6 ADVERSE REACTIONS

The following serious or otherwise important adverse reactions for bowel preparations are described elsewhere in the labeling:

- Serious Fluid and Electrolyte Abnormalities [see Warnings and Precautions (5.1)]
- Cardiac Arrhythmias [see Warnings and Precautions (5.2)]
- Seizures [see Warnings and Precautions (5.3)]
- Patients with Risk of Renal Injury [see Warnings and Precautions (5.4)]

^a Mild: barely noticeable, does not influence functioning causing no limitations of usual activities;

Moderate: makes participant uncomfortable, influences functioning causing some limitations of usual activities;

Severe: severe discomfort, treatment needed, severe and undesirable, causing inability to carry out usual activities

^b Study 2 was not designed to support comparative claims for SUTAB for the adverse reactions reported in this table.

^c Percentage represents n/N for patients who experienced each gastrointestinal adverse reaction on the symptom questionnaire based on the total number of patients per treatment arm.

Additional Adverse Reactions Reported in Studies 1 and 2

In addition to the gastrointestinal symptoms reported on the standard questionnaire (Tables 1 and 2), other adverse reactions reported in at least 2% of patients in either treatment arm in Studies 1 and 2 were: dizziness in Study 1 (0% SUTAB and 2% comparator); and hypermagnesemia (2% SUTAB and 2% comparator) and increased liver function test (including ALT, AST and bilirubin) (3% SUTAB and 1% comparator) in Study 2.

Laboratory Changes

Electrolyte Abnormalities

Shifts in serum electrolytes from normal at baseline to above the upper end of normal following study drug on the day of colonoscopy in at least 2% of patients in either treatment arm and at least 2% greater in patients treated with SUTAB than treated with comparator in either Study 1 or Study 2 were: magnesium (27% SUTAB and 5% comparator in Study 1), and serum osmolality (44% SUTAB and 28% comparator in Study 2). These changes were transient and resolved without intervention.

Renal Function Parameters

Decreases in creatinine clearance and increases in blood urea nitrogen (BUN) were reported in less than 1% of patients in both SUTAB and comparator arms in both trials.

6.2 Postmarketing Experience

The following adverse reactions have been identified during post-approval use of SUTAB. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Gastrointestinal: Gastric erythema, gastric ulceration, gastritis

General: Hypersensitivity reactions including rash, urticaria, pruritus, facial swelling, dyspnea and anaphylactic reaction.

7 DRUG INTERACTIONS

7.1 Drugs That May Increase Risks of Fluid and Electrolyte Abnormalities

Use caution when prescribing SUTAB to patients taking medications that increase the risk of fluid and electrolyte disturbances or may increase the risk of adverse events of seizure, arrhythmias, and prolonged QT in the setting of fluid and electrolyte abnormalities [see Warnings and Precautions (5.1, 5.2, 5.3, 5.4)].

7.2 Potential for Reduced Drug Absorption

SUTAB can reduce the absorption of other co-administered drugs [see Dosage and Administration (2.1)]:

- Administer oral medications at least one hour before starting each dose of SUTAB.
- Administer tetracycline and fluoroquinolone antibiotics, iron, digoxin, chlorpromazine, and penicillamine at least 2 hours before and not less than 6 hours after administration of each dose of SUTAB to avoid chelation with magnesium.

7.3 Stimulant Laxatives

Concurrent use of stimulant laxatives and SUTAB may increase the risk of mucosal ulceration or ischemic colitis. Avoid use of stimulant laxatives (e.g., bisacodyl, sodium picosulfate) while taking SUTAB [see Warnings and Precautions (5.5)].

USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

8

Risk Summary

There are no available data on SUTAB use in pregnant women to evaluate for a drug-associated risk of major birth defects, miscarriage, or adverse maternal or fetal outcomes. No reproduction or developmental studies in animals have been conducted with sodium sulfate, magnesium sulfate, and potassium chloride (SUTAB).

The estimated background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2% to 4% and 15% to 20%, respectively.

8.2 Lactation

Risk Summary

There are no available data on the presence of SUTAB in human or animal milk, the effects of on the breastfed child, or the effects on milk production.

The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for SUTAB and any potential adverse effects on the breastfed child from SUTAB or from the underlying maternal condition.

8.4 Pediatric Use

Safety and effectiveness in pediatric patients have not been established

8.5 Geriatric Use

Of the 471 patients who received SUTAB in pivotal clinical trials, 150 (32%) were 65 years of age or older, and 25 (5%) were 75 years of age or older. No differences in safety or effectiveness of SUTAB were observed between geriatric patients and younger patients. Elderly patients are more likely to have decreased hepatic, renal or cardiac function and may be more susceptible to adverse reactions resulting from fluid and electrolyte abnormalities [see Warnings and Precautions (5.1)].

8.6 Renal Impairment

Use SUTAB with caution in patients with renal impairment or patients taking concomitant medications that may affect renal function. These patients may be at risk for renal injury. Advise these patients of the importance of adequate hydration before, during and after use of SUTAB and consider performing baseline and post-colonoscopy laboratory tests (electrolytes, creatinine, and BUN) in these patients [see Warnings and Precautions (5.4)].

10 OVERDOSAGE

Overdosage of more than the recommended dose of SUTAB may lead to severe electrolyte disturbances, as well as dehydration and hypovolemia, with signs and symptoms of these disturbances [see Warnings and Precautions (5.1, 5.2, 5.3)]. Monitor for fluid and electrolyte disturbances and treat symptomatically.

11 DESCRIPTION

SUTAB (sodium sulfate, magnesium sulfate, and potassium chloride) tablets is an orally administered osmotic laxative and is provided as two bottles, each containing 12 tablets. Each tablet contains: 1.479 g sodium sulfate, 0.225 g magnesium sulfate, and 0.188 g potassium chloride. Inactive ingredients include: polyethylene glycol 8000, sodium caprylate, and ethylene glycol and vinyl alcohol graft copolymer.

Sodium Sulfate, USP

The molecular formula is Na₂SO₄. The average molecular weight is 142.04. The structural formula is:

Magnesium Sulfate, USP

The molecular formula is MgSO₄. The average molecular weight is 120.37. The structural formula is:

Potassium Chloride, USP

The molecular formula is KCI. The average molecular weight is 74.55. The structural formula is:

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

dogs up to 28 days up to a maximum daily dose of 5 grams/kg/day (approximately 0.9 and 3 times for rats and dogs, respectively, the recommended SUTAB human dose of 45.4 grams/day or 0.86 grams/kg based on the body surface area). In rats, the sulfate salts caused diarrhea and electrolyte and metabolic changes, including hypochloremia, hypokalemia, hyponatremia, lower serum osmolality, and high serum bicarbonate. Significant renal changes included increased fractional sodium excretion, increased urinary sodium and potassium excretion, and alkaline urine in both male and females. In addition, creatinine clearance was significantly decreased in females at the highest dose. No microscopic renal changes were seen. In dogs, the sulfate salts caused emesis, excessive salivation, excessive drinking of water, and abnormal excreta (soft and/or mucoid feces and/or diarrhea) and increased urine pH and sodium excretion.

14 CLINICAL STUDIES

The colon cleansing efficacy of SUTAB was evaluated in two randomized, single-blind, active-controlled, multicenter trials (Study 1 and Study 2). These trials included adult subjects undergoing colonoscopy for colorectal cancer screening and surveillance, or diagnostic colonoscopy, including subjects with abdominal pain, diarrhea, constipation and non-severe inflammatory bowel disease.

In Study 1 (BLI4700-301; NCT 03404401), 548 adult patients were included in the efficacy analysis. Patients ranged in age from 19 to 84 years (median age 59 years) and 56% were female. Racial distribution was 78% Caucasian, 16% African-American, and 11% Hispanic or Latino. Patients were randomized to one of the following two colon preparation regimens: SUTAB or polyethylene glycol 3350, sodium sulfate, sodium chloride, potassium chloride, ascorbic acid and sodium ascorbate for oral solution. Both preparations were administered according to a split-dose regimen [see Dosage and Administration (2.2)]. Patients receiving SUTAB were limited to a low residue breakfast followed by clear liquids on the day prior to the day of colonoscopy; patients receiving the comparator bowel prep were allowed to have a normal breakfast and a light lunch, followed by clear liquids and/ or yogurt for dinner. Approximately 97% of patients in the study completed both doses of preparation (98% of SUTAB patients and 95% of comparator patients).

In Study 2 (BLI4700-302; NCT 03261960), 388 adult patients were included in the efficacy analysis. Patients ranged in age from 23 to 83 years (median age 58 years) and 58% were female. Racial distribution was 94% Caucasian, 9% Hispanic or Latino, and 5% African-American. Patients were randomized to one of the following two colon preparation regimens: SUTAB or sodium picosulfate, magnesium oxide, and anhydrous citric acid for oral solution. Both preparations were administered according to a split-dose regimen [see Dosage and Administration (2.2)]. Patients receiving SUTAB were limited to a low residue breakfast followed by clear liquids on the day prior to the day of colonoscopy; patients receiving the comparator bowel prep were only allowed clear liquids on the day prior to colonoscopy. Approximately 98% of patients in the study completed both doses of preparation (98% of SUTAB patients and 99% of comparator patients).

The primary efficacy endpoint in each trial was the proportion of patients with successful colon cleansing, as assessed by the blinded colonoscopist utilizing the four-point scaled described below. Success was defined as an overall cleansing assessment of 3 (Good) or 4 (Excellent).

Score	Grade	Description
1	Poor	Large amount of fecal residue, additional bowel preparation required.
2	Fair	Enough feces even after washing and suctioning to prevent clear visualization of the entire colonic mucosa.
3	Good	Feces and fluid requiring washing and suctioning, but still achieves clear visualization of the entire colonic mucosa.
4	Excellent	No more than small bits of feces/fluid which can be suctioned easily; achieves clear visualization of the entire colonic mucosa.

Results for the primary endpoint in Studies 1 and 2 are shown in Table 3. In both trials, SUTAB was non-inferior to the comparator.

Table 3: Proportion of Adult Patients with Overall Cleansing Success^a in Two Controlled Trials with a Split-Dose Regimen

	SUTAB	% (n/N)	SUTAB	-comparator
	% (n/N)		Difference ^b (%)	99% Confidence Interval ^b
Study 1	92% (257/278)	89% ^c (241/270)	3.0	(-3.2, 9.3) ^e
Study 2	92% (175/190)	88% ^d (174/198)	3.1	(-4.5, 10.7) ^e

^a Success was defined as an overall cleaning assessment of 3 (Good) or 4 (Excellent) by the blinded endoscopist, scores were assigned on withdrawal of colonoscope.

⁹ treatment differences and confidence intervals were adjusted by study sites based on Mantel-Haenszel method comparator in Study 1 was polyethylene glycol 3350, sodium sulfate, sodium chloride, potassium chloride, sodium ascorbate and ascorbic acid for oral solution

⁴ comparator in Study 2 was sodium picosulfate, magnesium oxide, and anhydrous citric acid for oral solution e non-inferior

16 HOW SUPPLIED/STORAGE AND HANDLING

Each tablet of SUTAB contains 1.479 g sodium sulfate, 0.225 g magnesium sulfate, and 0.188 g potassium chloride. The tablets are white to off-white, film coated, oblong, and biconvex with flat sides, debossed with S24 on one side

Each carton of SUTAB (NDC 52268-201-01) contains:

- Two bottles, each bottle (NDC 52268-200-01) contains 12 tablets.
- One container with a 16-ounce fill line.

The primary mode of action is osmotic action of sodium sulfate and magnesium sulfate, which induce a laxative effect. The physiological consequence is increased water retention in the lumen of the colon, resulting in loose stools.

12.3 Pharmacokinetics

Absorption

After the oral administration of SUTAB to patients in clinical studies, the median serum sulfate concentration increased by about 2.5-fold at 5 to 8 hours post Dose 2 (0.61 mmol/L) compared to baseline (0.25 mmol/L) and returned to baseline by 24 to 48 hours after colonoscopy.

Elimination

Fecal excretion is the primary route of sulfate elimination.

Use in Specific Populations

Patients with Renal Impairment

The disposition of sulfate after ingestion of a sulfate-based product containing sodium sulfate, potassium sulfate, and magnesium sulfate similar to SUTAB was studied in patients (N=6) with moderate renal impairment (creatinine clearance of 30 to 49 mL/min). In patients with moderate renal impairment, mean AUC was 54% higher and mean C_{max} was 44% higher than healthy subjects. The mean sulfate concentrations in healthy subjects and in patients with moderate renal impairment returned to their respective baselines by Day 6 after dose initiation. Urinary excretion of sulfate over 30 hours after the first dose was approximately 16% lower in patients with moderate renal impairment than in healthy subjects. These differences are not considered clinically meaningful.

Patients with Hepatic Impairment

The disposition of sulfate after ingestion of a sulfate-based product containing sodium sulfate, potassium sulfate, and magnesium sulfate similar to SUTAB was also studied in patients (N=6) with mild-moderate hepatic impairment (Child-Pugh grades A and B). Systemic exposure of serum sulfate (AUC and C_{max}) was similar between healthy subjects and patients with hepatic impairment. The mean sulfate concentrations in healthy subjects and in patients with mild to moderate hepatic impairment returned to their respective baselines by Day 6 after dose initiation. Urinary excretion of sulfate over 30 hours after the first dose was similar between patients with hepatic impairment and healthy subjects.

13 NONCLINICAL TOXICOLOGY

13.2 Animal Toxicology and/or Pharmacology

Animal toxicology studies with sodium sulfate, magnesium sulfate, and potassium chloride (SUTAB) have not been conducted. Sulfate salts of sodium, potassium, and magnesium, were administered orally (gavage) to rats and

Storage

Store at 20° to 25°C (68° to 77°F). Excursions permitted between 15° to 30°C (59° to 86°F). See USP controlled room temperature

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Medication Guide and Instructions for Use). Instruct patients:

- Administration of two doses of SUTAB (24 tablets) are required for a complete preparation for colonoscopy. Twelve (12) tablets are equivalent to one dose.
- Water must be consumed with each dose of SUTAB and additonal water must be consumed after each dose.
- If preparation-related symptoms occur (e.g. nausea, bloating, cramping), to pause or slow the rate of drinking the additional water until symptoms diminish.
- Not to take other laxatives while they are taking SUTAB.
- Not to drink milk or eat or drink anything colored red or purple.
- Not to drink alcohol •
- Not to take oral medications within one hour of starting each dose of SUTAB. .
- If taking tetracycline or fluoroquinolone antibiotics, iron, digoxin, chlorpromazine, or penicillamine, take these • medications at least 2 hours before and not less than 6 hours after administration of each dose of SUTAB.
- To complete all SUTAB tablets and required water at least two hours prior to colonoscopy.
- To contact their healthcare provider if they develop significant vomiting or signs of dehydration after taking SUTAB or if they experience cardiac arrhythmias or seizures [see Warnings and Precautions (5.1, 5.2, 5.3)].

Manufactured by: Braintree Laboratories, Inc. 270 Centre Street Holbrook, MA 02343

This product is covered by the following patents: U.S. Patents 10,143,656, 11,033,498

Please see www.sebelapharma.com for patent information.

© Braintree Laboratories. Inc.

	1
Medication Guide	How should I take SUTAB?
SUTAB® (Sootab)	See the Instructions for Use for dosing instructions. You must read, understand, and
(sodium sulfate, magnesium sulfate, and potassium chloride)	follow these instructions to take SUTĂB the right way.
tablets, for oral use	Take SUTAB exactly as your healthcare provider tells you to take it.
Read and understand these Medication Guide Instructions at least 2 days before your	 Each dose of SUTAB is 12 tablets that must be taken while drinking 16 ounces of water. Two doses of SUTAB (24 tablets) are required for a complete colonoscopy preparation.
colonoscopy and again before you start taking SUTAB.	It is important for you to drink the additional prescribed amount of water listed in the
What is the most important information I should know about SUTAB?	Instructions for Use to prevent fluid loss (dehydration).
SUTAB and other bowel preparations can cause serious side effects, including:	• SUTAB is taken using the Split-Dose method. See the Instructions for Use for more
Serious loss of body fluid (dehydration) and changes in blood salts (electrolytes) in your	information.
blood.	 All people taking SUTAB should follow these general instructions starting 1 day before your colonoscopy:
These changes can cause:	 you can eat a low-residue breakfast. Low-residue foods include eggs, white bread,
abnormal heartbeats that can cause death.	cottage cheese, yogurt, grits, coffee and tea.
• seizures. This can happen even if you have never had a seizure.	\circ after breakfast only drink clear liquids all day and the next day until 2 hours before your
kidney problems.	colonoscopy. Stop drinking all fluids at least 2 hours before the colonoscopy.
Your chance of having fluid loss and changes in body salts with SUTAB is higher if you:	 after taking the first dose of SUTAB (12 tablets) if you have any bloating or feeling like your stomach is upset, wait to take your second dose of 12 tablets until your stomach
have heart problems.	feels better.
have kidney problems.	While taking SUTAB, do not:
• take water pills (diuretics) or non-steroidal anti-inflammatory drugs (NSAIDs).	 ○ take any other laxatives
Tell your healthcare provider right away if you have any of these symptoms of a loss of too much body fluid (dehydration) while taking SUTAB:	$^{\circ}$ take any medicines by mouth (oral) within 1 hour of starting SUTAB
vomiting • urinating less often than normal	$^{\circ}$ eat solid foods, dairy such as milk, or drink alcohol
dizziness headache	\circ eat or drink anything colored red or purple
See "What are the possible side effects of SUTAB?" for more information about side	Contact your healthcare provider right away if after taking SUTAB you have severe
effects.	vomiting, signs of dehydration, abnormal heartbeats or seizures.
What is SUTAB?	What are the possible side effects of SUTAB?
SUTAB is a prescription medicine used by adults to clean the colon before a colonoscopy.	SUTAB can cause serious side effects, including:
SUTAB cleans your colon by causing you to have diarrhea. Cleaning your colon helps your	• See "What is the most important information I should know about SUTAB?"
healthcare provider see the inside of your colon more clearly during your colonoscopy.	• Changes in certain blood tests. Your healthcare provider may do blood tests after you take
It is not known if SUTAB is safe and effective in children.	SUTAB to check your blood for changes. Tell your healthcare provider if you have any symptoms of too much fluid loss, including:
Do not take SUTAB if your healthcare provider has told you that you have:	 vomiting o nausea o bloating
 a blockage in your bowel (obstruction) or a problem with food moving too slowly through your intestines (ileus) 	 dizziness stomach (abdominal) cramping headache
 an opening in the wall of your stomach or intestine (bowel perforation) 	 ○ urinate less than usual
 a very dilated intestine (toxic colitis or toxic megacolon) 	 o trouble drinking clear liquid
 problems with food and fluid emptying from your stomach (gastric retention) 	Heart problems. SUTAB may cause abnormal heartbeats.
 allergies to any ingredient in SUTAB 	• Seizures.
	• Ulcers of the bowel or bowel problems (ischemic colitis). Tell your healthcare provider right away if you have severe stomach-area (abdominal) pain or rectal bleeding.
Before taking SUTAB, tell your healthcare provider about all of your medical conditions, including if you:	The most common side effects of SUTAB include:
 have heart problems. 	nausea vomiting
 have stomach or bowel problems including ulcerative colitis. 	 stomach-area swelling (abdominal distention) upper abdominal pain
 have problems with swallowing or gastric reflux. 	 stomach irritation
 have a history of seizures. 	These are not all the possible side effects of SUTAB.
• are withdrawing from drinking alcohol or from taking benzodiazepines.	Call your doctor for medical advice about side effects. You may report side effects to FDA at
have a low blood salt (sodium) level.	1-800- FDA-1088.
have kidney problems.	How should I store SUTAB?
• are pregnant. It is not known if SUTAB will harm your unborn baby. Talk to your healthcare	Store SUTAB at room temperature between 68°F to 77°F (20°C to 25°C).
provider if you are pregnant.	Keep SUTAB and all medicines out of the reach of children.
are breastfeeding or plan to breastfeed. It is not known if SUTAB passes into your breast milk. You and your bealthcare provider should decide if you will take SUTAB while	General information about the safe and effective use of SUTAB.
milk. You and your healthcare provider should decide if you will take SUTAB while breastfeeding.	Medicines are sometimes prescribed for purposes other than those listed in a Medication
Tell your healthcare provider about all the medicines you take, including prescription and	Guide. Do not use SUTAB for a condition for which it was not prescribed. Do not give SUTAB to other people, even if they are going to have the same procedure you are. It may harm them.
over-the-counter medicines, vitamins, and herbal supplements.	You can ask your pharmacist or healthcare provider for information that is written for health
SUTAB may affect how other medicines work. Medicines taken by mouth may not be absorbed	professionals.
properly when taken within 1 hour before the start of each dose of SUTAB.	What are the ingredients in SUTAB?
Especially tell your healthcare provider if you take:	Active ingredients: sodium sulfate, magnesium sulfate, and potassium chloride
medicines for blood pressure or heart problems. medicines for kidney problems	Inactive ingredients: polyethylene glycol 8000, sodium caprylate, and ethylene glycol and
medicines for kidney problems. medicines for solution	vinyl alcohol graft copolymer
medicines for seizures.	
water pills (diuretics). pop-steroidal anti-inflammatory drugs (NSAIDs)	
 non-steroidal anti-inflammatory drugs (NSAIDs). medicines for depression or other montal bactth problems 	
 medicines for depression or other mental health problems. laxatives. Do not take other laxatives while taking SUTAB. 	
The following medicines should be taken at least 2 hours before starting each dose of SUTAB	
and not less than 6 hours after taking each dose of SUTAB:	
tetracycline iron chlorpromazine	
fluoroquinolone antibiotics	Manufactured by:
Ask your healthcare provider or pharmacist for a list of these medicines if you are not sure if	Braintree Laboratories, Inc.
you are taking any of the medicines listed above.	270 Centre Street Holbrook, MA 02343
Know the medicines you take. Keep a list of them to show your healthcare provider and	For more information, go to <u>www.braintreelabs.com</u> or call 1-800-874-6756.
pharmacist when you get a new medicine.	To more mornation, yo to <u>www.brainticelabs.com</u> of call 1-000-0/4-0/30.